

DISCLOSURE: The following authors disclosed financial relationships: K. Waki received lecture fees from Takeda Pharmaceutical. S. Shichijo received lecture fees from Olympus, Boston Scientific Japan, Daiichi-Sankyo, EA Pharma, Zeria Pharmaceutical, The Japanese Society of Gastroenterology, and the Japan Gastroenterological Endoscopy Society. N. Uedo received fees for lectures, presentations, and educational events from Olympus, FUJIFILM, Boston Scientific Japan, Daiichi-Sankyo, Takeda Pharmaceutical, EA Pharma, Otsuka Pharmaceutical, and AstraZeneca. Y. Takeuchi received consultant fees from the Pharmaceuticals and Medical Devices Agency and EPARK, Inc., as well as lecture fees from SAS Institute Japan Ltd. Additionally, Y. Takeuchi is involved in a collaborative study with Pfizer Inc., which is not related to this article. T. Kanesaka received lecture fees from Olympus. Y. Takeuchi received lecture fees from Olympus, Boston Scientific Japan, Daiichi-Sankyo, Miyarisan Pharmaceutical, AstraZeneca, EA Pharma, Zeria Pharmaceutical, FUJIFILM, Kaneka Medix, Kyorin Pharmaceutical, and the Japan Gastroenterological Endoscopy Society. All other authors disclosed no financial relationships.