

DISCLOSURE: The following authors disclosed financial relationships: S. Shichijo: Speaker for Olympus, Boston Scientific Japan, Daiichi-Sankyo, EA Pharma, Zeria Pharmaceutical, The Japanese Society of Gastroenterology, and Japan Gastroenterological Endoscopy Society. Y. Takeuchi: Speaker for Olympus, Boston Scientific Japan, Daiichi- Sankyo, Miyarisan Pharmaceutical, Asuka Pharmaceutical, AstraZeneca, EA Pharma, Zeria Pharmaceutical, Fujifilm, Kaneka Medix, Kyorin Pharmaceutical, and Japan Gastroenterological Endoscopy Society. Y. Sano: Speaker for Olympus, AbbVie, Ono Pharmaceutical, EA Pharma, Boston Scientific Japan, Japan Gastroenterological Endoscopy Society, Takeda Pharmaceutical, Otsuka Pharmaceutical, AstraZeneca, Torii Pharmaceutical, Chugai Pharmaceutical, and Mylan EPD; technical guidance fee from Olympus; royalties from Nankodo, Medical View, and Nihon Medical Center. M. Kitano: Honoraria from Olympus; grant recipient from Zeon Medical, Zeon Japan, and Boston Scientific Japan. All other authors disclosed no financial relationships. Research support for this study was provided by the Kansai Endoscopic Device Selection Conference in the Kansai Research Group. The funder paid for the data center that constructed the data allocation and collecting system.